

Chronic Maintenance In-Center Hemodialysis Standing Orders - Iron

Sodium ferric gluconate (ICD10- D63.1)

- **1. Goal**: Iron saturation 30 50%; Ferritin 500 1200 ng/ml.
- **2. Labs:** (ICD10 = E83.10)
 - a. Draw iron labs quarterly (ferritin and iron saturation) Jan.-Apr.-Jul.-Oct. or as indicated by table below.
 - b. Wait a minimum of 7 days after the last dose of Sodium ferric gluconate before drawing iron labs. May wait until next monthly lab draw.
- **3. Test Dose**: Administer Sodium ferric gluconate test dose of 25 mg (2 cc of a 62.5 mg vial) IV over 3 minutes for the first dose only to assure no allergic reaction.
- **4. Infection/Antibiotics:** HOLD IV iron if patient has signs of significant infection or is on antibiotics.

5. Sodium ferric gluconate dosing:

- a. Based on most recent iron labs give IV Sodium ferric gluconate and draw labs per table below.
- b. Doses with frequency "q run" may be given up to 3x/week depending on patient's dialysis schedule.
- c. Weekly dosing should be given on the mid-week run.

If	And	Actions		
Ferritin Iron < 500 Saturation		IV Sodium ferric gluconate Dose	Draw Iron Labs	
	< 20%	125 mg q run ×9	7 days after last dose Based on labs move to new dose	
	20 - 30%	125 mg q run × 6	7 days after last dose Based on labs move to new dose 7 days after last dose Based on labs move to new dose te Quarterly	
	31 - 50%	125 mg q week x 3		
	> 50%	Hold Sodium ferric gluconate		

Patient Name	NKC#
MEC approved 12.13.18	Page 1 of 2

Northwest Kidney Centers

Chronic Maintenance In-Center Hemodialysis Standing Orders – Iron

If	And	Actions		
Ferritin 500 - 800	Iron Saturation	IV Sodium ferric gluconate Dose	Draw Iron Labs	
	< 20%	125 mg q run × 6	7 days after last dose Based on labs move to new dose	
20 - 30%		125 mg q run x 3 then 62.5 mg q 2 weeks	Quarterly	
	31 - 50%	62.5 mg q week x 3 then 62.5 mg q month	Quarterly	
	>50%	Hold Sodium ferric gluconate	Quarterly	
If	And	Actions		
Ferritin 801 - 1200	Iron Saturation	IV Sodium ferric gluconate Dose	Draw Iron Labs	
	< 20%	62.5 mg q 2 weeks x 2	7 days after last dose Check with MD re: further doses	
	20 - 30%	62.5 mg q month x 2	Quarterly	
	31 - 50%	Hold Sodium ferric gluconate	Quarterly	
	>50%	Hold Sodium ferric gluconate	Quarterly	
If	And	Actions		
Ferritin >1200			Draw Iron Labs	
	< 20 %	Check with MD	Quarterly (unless otherwise ordered)	
	20 - 30 %	Hold Sodium ferric gluconate	Quarterly	
	31 - 50 %	Hold Sodium ferric gluconate	Quarterly	
	> 50 %	Hold Sodium ferric gluconate	Quarterly	

6. Hemoglobin:

- a. If hemoglobin \geq 12, iron saturation > 30%, and ferritin > 800, hold Sodium ferric gluconate.
- b. If hemoglobin \geq 12, iron saturation \leq 30%, contact MD for direction.

Physician Name (Please Print)	RN Name (Please Print)	
Physician signature (see Initial Orders)	RN signature	Date
Patient Name	NKC#	