

Chronic Maintenance In-Center Hemodialysis Standing Orders - Iron

Sodium ferric gluconate (ICD10- D63.1)

1. Goal: Iron saturation 30 - 50%

2. Labs: (ICD10 = E83.10)

- Draw iron labs quarterly (ferritin and iron saturation) Jan.-Apr.-Jul.-Oct.
- Wait a minimum of 7 days after the last dose of Sodium ferric gluconate before drawing iron labs.

3. Hemoglobin:

- If hemoglobin is greater than or equal to 12 g/dL or ferritin is greater than or equal to 1000 ng/mL, hold Sodium ferric gluconate.
- When hemoglobin then decreases to less than 12 g/dL, restart iron per protocol.

4. Infection/Antibiotics: HOLD IV iron if patient has an infection requiring IV antibiotics. If patient is still receiving IV antibiotics more than 2 weeks, then can resume IV iron dosing at that time.

5. Sodium ferric gluconate dosing:

- Based on most recent iron labs, give IV Sodium ferric gluconate per table below.
- Doses with frequency "every run" may be given up to 3x/week depending on patient's dialysis schedule.
- Maintenance dosing: 125 mg given every 4 weeks. Dose should start the second full week of the month (mid-week run preferable).

If	Actions
Iron Saturation	IV Sodium ferric gluconate Dose
< 20%	125 mg every run ×6, then give monthly maintenance dose the following month
20 - < 30%	125 mg every run ×3, then give monthly maintenance dose the following month
30 - 50%	125 mg every month (maintenance dose)
> 50%	Hold Sodium ferric gluconate, recheck iron saturation every month <i>until iron saturation falls below 50% again.</i> Follow rules above.

Matthew Rivara, MD

Physician Name (Please Print)

Physician signature

(see **Initial Orders**)

Date

Patient Name

NKC#